


AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the Application.

5 Listing of Claims

1. (Currently Amended) A compound of the formula:



wherein:

a) one of R¹ and R³ is H and R², R⁴; and the other of R¹ and R³ are independently selected from H and a group of the formula:




wherein:


i) n is an integer from 1 to 12;

ii) each L is independently selected from a member of the group D1, D2, E, or J wherein:

D1 is of the formula:




D2 is of the formula:




E is of the formula:

4




J is of the formula:



wherein:

A) each X and Y is independently a group of the formula:



and

B) m is an integer from 0 to 8; and

C) no more than two D2 or E groups are adjacent to each other and no J groups are adjacent to each other;

iii) each Z is independently selected from:

A) hydrogen, halo, cyano, nitro, nitroso, azido, chlorate, bromate, iodate, isocyanato, isocyanido, isothiocyanato, pentafluorothio, or

B) a group G, wherein G is an unsubstituted or substituted; unsaturated, partially saturated, or saturated; monocyclic, bicyclic, tricyclic, or fused; 4 to 14 membered carbocyclic or heterocyclic ring system wherein;

1) when the ring system contains a 4 membered heterocyclic ring, the heterocyclic ring contains 1 heteroatom;

5


- 2) when the ring system contains a 5, or more, membered heterocyclic ring or a polycyclic heterocyclic ring, the heterocyclic or polycyclic heterocyclic ring contains from 1 to 4 heteroatoms;
- 3) each heteroatom is independently selected from N, O, and S;
- 5 4) the number of substituents is from 0 to 5 and each substituent is independently selected from X;
- b) the total number of non-hydrogen atoms in each compound is 50 or less; and
- c) the total number of heteroatoms in $-(L)_n-Z$ is from 0 to 4; and
- d) either;
- 10 i) R^1 or R^3 contains at least one group G; or
- ii) at least one L group is an E group; or
- iii) at least one of R^1 , R^2 , R^3 , and R^4 contains one to four non-hydrogen atoms and at least one of R^1 , R^2 , R^3 , and R^4 contains more than four non-hydrogen atoms; _____

15 and its enantiomers, stereoisomers, salts, and mixtures thereof;
or a composition thereof;

provided that:

- a) $-(L)_n-Z$ is other than trimethylsilyl, trimethylsilylsulfonyl or thiol; and
- b) R^1 is other than phenylsulfonyl, phenylthioethyl, diphenylhydroxymethyl, benzo[g]quinolin-7-yl-1-methyl, a malonate derivative, a substituted 3-aminocyclohexenone, a dialkoxybenzylaminocarbonyl; and
- 20 c) R^3 is other than 2-phenyl-ethenyl, phenylthio, (4-bromo-2-methylphenyl)carbamic acid N-carbonyl, (4-bromo-2-methylphenyl)carbamic acid ethyl ester N-carbonyl, a malonate derivative, aryloxy, or a
- 25 dialkoxybenzylaminecarbonyl.

2. (Currently Amended) A method of inhibiting an ethylene response in a plant comprising the step of contacting the plant with an effective ethylene response-inhibiting amount of a cyclopropene derivative of the formula:



6

wherein:

- a) one of R^1 and R^3 is H and R^2, R^4 ; and the other of R^1 and R^3 are independently selected from H and a group of the formula:




5


wherein:

- i) n is an integer from $[[1]]$ 0 to 12;
 ii) each L is independently selected from a member of the group D1, D2, E, or J wherein:

D1 is of the formula:




D2 is of the formula:




15

E is of the formula:




J is of the formula:

7



wherein:

A) each X and Y is independently a group of the formula:



and

B) m is an integer from 0 to 8; and

C) no more than two D2 or E groups are adjacent to each other and no J groups are adjacent to each other;

iii) each Z is independently selected from:

A) hydrogen, halo, cyano, nitro, nitroso, azido, chlorate, bromate, iodate, isocyanato, isocyanido, isothiocyanato, pentafluorothio, or

B) a group G, wherein G is an unsubstituted or substituted; unsaturated, partially saturated, or saturated; monocyclic, bicyclic, tricyclic, or fused; $[[4]]$ 3 to 14 membered carbocyclic or heterocyclic ring system wherein;

1) when the ring system contains a 3 or 4 membered heterocyclic ring, the heterocyclic ring contains 1 heteroatom;

2) when the ring system contains a 5, or more, membered heterocyclic ring or a polycyclic heterocyclic ring, the heterocyclic or polycyclic heterocyclic ring contains from 1 to 4 heteroatoms;

3) each heteroatom is independently selected from N, O, and S;

4) the number of substituents is from 0 to 5 and each substituent is independently selected from X;

b) the total number of non-hydrogen atoms in each compound is 50 or less; and

c) the total number of heteroatoms in $-(\text{L})_m-\text{Z}$ is from 0 to 4; and

d) either;

i) R^1 or R^3 contains at least one group G; or

ii) at least one L group is an E group; or

iii) at least one of R^1 , R^2 , R^3 , and R^4 contains one to four non-hydrogen atoms and at least one of R^1 , R^2 , R^3 , and R^4 contains more than four non-hydrogen atoms; _____

and its enantiomers, stereoisomers, salts, and mixtures thereof;

5 or a composition thereof;

3. (Original) The method of claim 2, wherein the ethylene response is one or more of ripening or senescence of flowers, fruits, and vegetables; abscission of foliage, flowers, and fruit; the shortening of life of ornamental plants, cut flowers, shrubbery, seeds, or dormant seedlings; inhibition of growth; stimulation of growth; auxin activity; inhibition of terminal growth; control of apical dominance; increase in branching; increase in tillering; changing the morphology of plants, modifying the susceptibility to plant pathogens such as fungi, changing bio-chemical compositions; abortion or inhibition of flowering or seed development; lodging effects; stimulation of seed germination; breaking of dormancy; hormone effects; and epinasty effects.
- 10 4. (Original) The method of claim 2, wherein R^2 , R^3 , and R^4 are hydrogen or R^1 , R^2 , and R^3 are hydrogen.
- 15 5. (Original) The method of claim 2, wherein n is from 1 to 7.
6. (Original) The method of claim 2, wherein m is from 0 to 2.
7. (Original) The method of claim 2, wherein:
- 20 a) each D1 is -CXY-, -CO-, or -CS-;
- b) each D2 is -NX- or -O-;
- c) each E is -S-, -SiXY-, or -SO₂-;
- d) each X and Y is independently H, halo, OH, SH, -C(O)(C₁-C₄)alkyl, -C(O)O(C₁-C₄)alkyl, -O-(C₁-C₄)alkyl, -S-(C₁-C₄)alkyl, or substituted or unsubstituted (C₁-C₄)alkyl; and
- 25 e) each Z is independently H, halo, or G.
8. (Original) The method of claim 2, wherein each G is independently a substituted or unsubstituted; five, six, or seven membered; aryl, heteroaryl, heterocyclyl, or cycloalkyl ring.
- 30 9. (Original) The method of claim 8, wherein each G is independently a substituted or unsubstituted phenyl, pyridyl, cyclohexyl, cyclopentyl, pyrrolyl, furyl, thiophenyl, triazolyl, pyrazolyl, 1,3-dioxolanyl, or morpholinyl.

9

10.(Original) The method of claim 8, wherein the substituents, when present, are independently selected from 1 to 3 of methyl, methoxy, and halo.

BEST AVAILABLE COPY